



## LACTATE DEHYDROGENASE (LDH) AND RESPONSE TO RENAL ARTERY STENTING.

ACC Poster Contributions

Ernest N. Morial Convention Center, Hall F

Sunday, April 03, 2011, 10:00 a.m.-11:15 a.m.

Session Title: Renal Dysfunction and Vascular Diseases

Abstract Category: 11. Peripheral Arterial/Carotid Disease/Aortic Disease

Session-Poster Board Number: 1009-104

Authors: Mohammad Khalil Kanjwal, Dong Zhang, Satjit Adlakha, Haifeng Yu, Pamela S. Brewster, Katherine Hoetzel, Steven Haller, Michael Steffes, Joseph Shapiro, William R. Colyer, Christopher J. Cooper, The University Of Toledo, Toledo, OH, University of Minnesota, Minneapolis, MN

**Background:** Lactic acid dehydrogenase (LDH) is a marker of cell injury or death that increases with renal infarction; however, the utility for detecting renal injury, before, during or after renal artery stenting (RAS) has not been studied.

**Methods:** One hundred patients undergoing RAS at 7 centers were randomized to embolic protection or double-blind use of a GPIIb/IIIa inhibitor in a 2x2 design. All data were analyzed blindly at independent core labs. GFR change was classified as improved (>20%), stable (-20% - +20%), or worsened (<-20%).

**Results:** There was a significant negative correlation between baseline LDH and baseline GFR (-0.242,  $p < 0.05$ ) that persisted at 24 hours as well as 1-month (Figure 1). Following RAS LDH decreased at 24-hours ( $190 \pm 58$  to  $177 \pm 50$ ,  $p < 0.05$ ) but increased at 1-month follow-up ( $213 \pm 60$ ). Improved GFR 1-month following RAS was associated with high LDH ( $p = 0.02$ ) (Figure 2). In contrast worsened GFR after RAS was associated with lower initial LDH values that rose significantly at 1-month follow-up ( $+36$ ,  $p = 0.006$ ) (Figure 2).

**Conclusions:** In atherosclerotic renal stenosis high baseline LDH is associated with low baseline GFR but improvement in GFR 1-month post procedure, potentially indicating ongoing renal injury amenable to treatment with stenting. In contrast, in patients with worsened GFR 1-month post procedure, LDH rises significantly suggesting ongoing injury late after the procedure.

